

Приложение 2

Характеристика групп пациентов на этапе финального осмотра (по данным литературы)

[illegible]

Jiang et al. [26]	Percutaneous group	–	–	–	–	–	–	–	–	–	$3,6 \pm 0,3$	$13,5 \pm 6,1$
Kim et al. [27]	–	–	20,6	–	–	–	–	–	–	–	2,2	–
Li et al. [28]	SSPI group	$7,5 \pm 5,2$	–	–	–	–	–	–	–	–	$1,1 \pm 0,6$	–
Wang et al. [29]	–	2,1	–	100,0	–	–	–	–	–	–	–	–
Zhang et al. [30]	–	$5,3 \pm 3,7$	$5,7 \pm 3,2$	–	–	–	–	–	–	–	$2,0 \pm 0,7$	$34,0 \pm 4,0$
Chou et al. [31]	Non-fusion group	$13,8 \pm 6,6$	–	100,0	–	–	–	–	–	–	$2,1 \pm 0,9$	–
Proietti et al. [32]	Group A	–	–	–	–	–	–	–	–	–	1,8	12,0
	Group B	–	–	–	–	–	–	–	–	–	4,3	38,0
Takami et al. [33]	–	-0,6	–	100,0	–	–	–	–	–	–	–	–
Vanek et al. [34]	MIS group	$4,4 \pm 9,4$	–	–	–	–	–	17	–	–	–	–
Zhao et al. [35]	PFFV group	–	–	–	29	3	0	–	–	–	–	–
	TSSF group	–	–	–	30	5	0	–	–	–	–	–
Fu et al. [36]	Opsf-4	–	–	100,0	–	–	–	–	–	–	–	–

[illegible]

Hoffman et al. [51]	CG group	–	–	–	–	–	–	–	–	–	–	$21,4 \pm 23,7$
	IG group	–	–	–	–	–	–	–	–	–	–	$17,7 \pm 11,8$
Perna et al. [52]	Group A	$8,7 \pm 4,8$	–	–	–	–	–	–	–	–	$4,5 \pm 1,8$	$27,3 \pm 10,1$
Zhu et al. [53]	MIS-F group	–	$6,1 \pm 5,1$	–	–	–	–	–	–	–	$1,2 \pm 0,5$	$11,5 \pm 2,3$
	MIS-O group	–	$7,9 \pm 10,1$	–	–	–	–	–	–	–	$1,2 \pm 0,8$	$12,0 \pm 2,1$
	Open-C group	–	$6,8 \pm 8,1$	–	–	–	–	–	–	–	$1,4 \pm 0,7$	$12,2 \pm 2,6$